

Accepted Manuscript

Regulation of arginine methyltransferase 3 by a *Wolbachia*-induced microRNA in *Aedes aegypti* and its effect on *Wolbachia* and dengue virus replication

Guangmei Zhang, Mazhar Hussain, Sassan Asgari



PII: S0965-1748(14)00135-0

DOI: [10.1016/j.ibmb.2014.08.003](https://doi.org/10.1016/j.ibmb.2014.08.003)

Reference: IB 2607

To appear in: *Insect Biochemistry and Molecular Biology*

Received Date: 26 June 2014


Revised Date: 9 August 2014

Accepted Date: 13 August 2014

Please cite this article as: Zhang, G., Hussain, M., Asgari, S., Regulation of arginine methyltransferase 3 by a *Wolbachia*-induced microRNA in *Aedes aegypti* and its effect on *Wolbachia* and dengue virus replication, *Insect Biochemistry and Molecular Biology* (2014), doi: 10.1016/j.ibmb.2014.08.003.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Graphical abstract



Regulation of arginine methyltransferase 3 by a *Wolbachia*-induced microRNA in *Aedes aegypti* and its effect on *Wolbachia* and dengue virus replication

Guangmei Zhang, Mazhar Hussain, and Sassan Asgari*

Australian Infectious Disease Research Centre, School of Biological Sciences, The University of Queensland, Brisbane, QLD 4072, Australia

*Corresponding author: Sassan Asgari, Australian Infectious Disease Research Centre, School of Biological Sciences, The University of Queensland, Brisbane Qld 4072, Australia. Tel: +617 3365 2043; Fax: +617 3365 1655; s.asgari@uq.edu.au

Abstract

The gram-negative endosymbiotic bacteria, *Wolbachia*, have been found to colonize a wide range of invertebrates, including over 40% of insect species. Best known for host reproductive manipulations, some strains of *Wolbachia* have been shown to reduce the host life span by about 50% and inhibit replication and transmission of dengue virus (DENV) in the mosquito vector, *Aedes aegypti*. The molecular mechanisms underlying these effects still are not well understood. Our previous studies showed that *Wolbachia* uses host microRNAs (miRNAs) to manipulate host gene expression for its efficient maintenance and limiting replication of DENV in *Ae. aegypti*. Protein arginine methyltransferases are structurally and functionally conserved proteins from yeast to human. In mammals, it has been reported that protein arginine methyltransferases such as PRMT1, 5 and 6 could regulate replication of different viruses. *Ae. aegypti* contains eight members of protein arginine methyltransferases (AaArgM1-8). Here, we show that the wMelPop strain of *Wolbachia* introduced into *Ae. aegypti* significantly induces the expression of AaArgM3. Interestingly, we found that *Wolbachia* uses aae-miR-2940, which is highly upregulated in *Wolbachia*-infected mosquitoes, to upregulate the expression of AaArgM3. Silencing of AaArgM3 in a mosquito cell line led to the inhibition of *Wolbachia* replication, but had no effect on the replication of DENV. These results provide further evidence that *Wolbachia* uses the host miRNAs to manipulate host gene expression and facilitate colonization in *Ae. aegypti* mosquito.

Key words: protein arginine methyltransferase 3; *Aedes aegypti*; *Wolbachia*; microRNA; dengue virus

1. Introduction

Wolbachia, the maternally inherited and gram-negative endosymbiotic bacteria, naturally occur in 40-65% of insect species (Hilgenboecker et al., 2008; Jeyapragash and Hoy, 2000; Zug and Hammerstein, 2012). In the absence of naturally present strains of *Wolbachia* in the main vectors of

dengue virus (DENV; *Aedes aegypti*) and malaria (*Anopheles gambiae*), *Wolbachia* strains from *Drosophila melanogaster* and *Ae. albopictus* have recently been successfully introduced into *Ae. aegypti* and other mosquito species (Bian et al., 2013; McMeniman et al., 2009; Xi et al., 2005). Although a recent study found natural infections of *Wolbachia* in *An. gambiae* field populations in Burkina Faso, West Africa (Baldini et al., 2014). In some cases, transinfected *Wolbachia* strains have established stable inherited infections in the lab and the field (Frentiu et al., 2014; Walker et al., 2011). Similar to their original hosts, the newly introduced *Wolbachia* strains induce cytoplasmic incompatibility and life-shortening in adult mosquitoes by as much as 50% (McMeniman et al., 2009; Moreira et al., 2009; Xi et al., 2005). In addition, *Ae. aegypti* infected with *Wolbachia* possesses very strong resistance to several arboviruses including DENV and Chikungunya virus (Bian et al., 2013; Moreira et al., 2009), and *Plasmodium* (Moreira et al., 2009) and filarial nematodes (Kambris et al., 2009). Thus, the utilization of *Wolbachia* to control arbovirus transmission from mosquitoes to vertebrate hosts has become one of the most exciting approaches in vector-borne disease control.

The molecular mechanism(s) underlying suppression of replication of viruses in the presence of *Wolbachia* are thought to be complex and perhaps due to a combination of factors, but still largely unknown (see a recent review (Rainey et al., 2014). In its natural host, *D. melanogaster*, *Wolbachia* confer host resistance to RNA viruses and other pathogens via non-immune related mechanisms, since *Wolbachia* did not induce expression of innate immune genes (Bourtzis et al., 2000; Rances et al., 2013; Rancès et al., 2012). In *Ae. aegypti*, studies have shown that *Wolbachia* could use innate immune related mechanisms to suppress the replication of DENV by inducing the production of reactive oxygen species (ROS), overexpression of host immune genes and production of a variety of antimicrobial effectors (Bian et al., 2010; Kambris et al., 2010; Kambris et al., 2009; Moreira et al., 2009; Pan et al., 2012; Xi et al., 2008). Recently, our studies demonstrated that *Wolbachia* use host microRNAs (miRNAs) to manipulate the expression of several host genes such as the metalloprotease ftsh, MCT, MCM6 and AaDnmt2, which facilitate *Wolbachia* colonization and

some contribute to inhibition of DENV replication in *Ae. aegypti* (Hussain et al., 2011; Osei-Amo et al., 2012; Zhang et al., 2013).

miRNAs are an evolutionarily conserved class of small non-coding RNAs (~22 nucleotides), which down- or upregulate gene expression via partial or complete complementarity to their target gene sequences. They play important roles in cellular processes including development, differentiation, apoptosis, immunity and host-microorganism interactions (reviewed in Asgari, 2013; Bartel, 2009). miRNAs may bind to the 3'UTR, 5'UTR or coding region of target genes. Previous studies have shown that one miRNA could target several genes or several miRNAs could target one gene (e.g. Osei-Amo et al., 2012; Zhang et al., 2013). The expression levels of cellular miRNAs may substantially change in response to bacterial and viral infections in animals and plants (Fehri et al., 2010; Huang et al., 2007; Hussain et al., 2011; Lu et al., 2008; Tili et al., 2007). In our previous studies, we found differential expression of several miRNAs in *Wolbachia*-infected *Ae. aegypti* mosquitoes (Hussain et al., 2011) leading to up- or downregulation of a variety of host genes, which facilitate colonization and host resistance to DENV in *Ae. aegypti* (Hussain et al., 2011; Osei-Amo et al., 2012; Zhang et al., 2013).

In this study, we identified protein arginine methyltransferase 3 (*AaArgM3*) as another target gene of the *Wolbachia*-induced mosquito-specific aae-miR-2940-5p in *Ae. aegypti*. *AaArgM3* belongs to protein arginine methyltransferase family, which includes eight members in *Ae. aegypti* (denoted *AaArgM1-8*). Arginine methyltransferases play diverse functions in cellular functions such as RNA processing and transcription (reviewed in Bedford and Clarke, 2009) and host-pathogen interactions (e.g. Duong et al., 2005; Souki et al., 2009; Yu et al., 2010). We investigated the effect of *Wolbachia* and DENV on these miRNAs and in turn their effect on replication of the two microorganisms. Our results suggest that *AaArgM3* plays an important role in the maintenance of *Wolbachia* infection in mosquito cells but has no effect on DENV replication.

2. Materials and Methods

2.1. Mosquitoes and insect cell lines

Ae. aegypti infected with the *wMelPop-CLA* strain of *Wolbachia* (*Wol*⁺) and a *Wolbachia*-free strain, tetracycline-cured line (*Wol*⁻), were the stocks as previously described (McMeniman et al., 2009). *Ae. aegypti* was reared at 25 °C with 80% relative humidity and a 12-h light regime. Larvae were maintained with fish food pellets (Tetramin, tetra) at a density of 50 larvae per litre water in flat trays. Adults were supplied 10% (W/V) sucrose solution, *ad libitum*. *Ae. aegypti* Aag2 cells and *wMelPop* infected Aag2 cells (denoted as aag2.*wMelPop-CLA*) (Frentiu et al., 2010) were maintained in a 1:1 mixture of Mitsunashi-Maramorosch and Schneider's insect media (Invitrogen) supplemented with 10% FBS.

2.2. RNA extraction, cDNA synthesis and polymerase chain reaction (PCR)

Total RNA from female and male mosquitoes (separately) and mosquito cell lines was isolated using Tri-Reagent (Molecular Research Center). The RNA was treated with DNase I before used for reverse transcription (RT). The first strand cDNA was synthesized by RT with a Poly(dT) primer. In each RT reaction, approximately 2 µg of total RNA was used as template in a total volume of 20 µl. Following cDNA synthesis, 2 µl of RT products were used for each PCR in a total reaction volume of 25 µl with *AaArgM3* gene-specific primers (Forward: 5'-GTAGACGTAGACTGTCCC-3'; Reverse: 5'-ACCGGAATCGGTTTCCTCG-3'). The amplification was performed at 94 °C for 3 min, followed by 35 cycles of 94 °C for 30 sec, 56 °C for 30 sec, 68 °C for 1 min, and a final extension at 68 °C for 5 min. The ribosomal protein S17 (*RPS17*) gene was used as control.

2.3. Quantitative PCR (qPCR) of *Wolbachia* density

Total genomic DNA was extracted from aag2.*wMelPop-CLA* cells. *Wolbachia* density in cells was determined by qPCR using the *wsp* gene-specific primers as described previously (Zhang et al., 2013). qPCR was carried out by using Platinum SYBR Green Mix (Invitrogen) with 20 ng of total genomic DNA in a Rotor-Gene thermal cycler (QIAGEN) under the following conditions: 95 °C

hold for 30 sec, then 40 cycles of 95 °C for 15 sec, 50 °C for 15 sec and 72 °C for 20 sec, followed by the melting curve analysis (68 °C to 95 °C). For this experiment, three biological replicates with three technical replicates were analysed. The *RPS17* gene was used for normalization of DNA templates. The student's *t* test was used to compare the differences in means between different treatments.

2.4. RT-qPCR

For RNA samples from mock and DENV-2 infected female mosquitoes, samples produced previously were utilized (Zhang et al., 2013). Following the RT reaction, qPCR with DENV gene-specific primers (forward: 5'-GTGGTGGTGACTGAGGACTG-3'; reverse: 5'-CCATCCCGTACCAGCATCCG-3') was carried out to determine DENV genomic RNA (gRNA) levels in cells. Platinum SYBR Green Mix (Invitrogen) was used for qPCR with 1 µl of RT products as described above. For this experiment, three biological replicates with three technical replicates were analysed. The *RPS17* gene was also used for normalization of RNA templates. The student's *t* test or ANOVA was used to compare the differences in means.

For tissue-specific analysis of *AaArgM3* transcript levels, total RNA was extracted from ovaries, salivary glands, thoracic muscle, midgut and fat body dissected from 4-day-old Wol⁺ and Wol⁻ female mosquitoes (Zhang et al., 2013). RT-qPCR reactions were performed using *AaArgM3* gene-specific primers as described above. Similarly, three biological replicates with three technical replicates were analysed for each tissue of mosquito type. Each biological replicate consisted of a pool of total RNA extracted from different tissues of 10 female mosquitoes. The *RPS17* gene was also used for normalization of RNA templates.

2.5. miRNA target prediction and validation

NCBI BLAST (<http://www.ncbi.nih.gov/BLAST>), RNAHybrid (Rehmsmeier et al., 2004) and RNA22 software (IBM) were used to identify the potential miRNAs induced in *Wolbachia*-infected female mosquitoes interacting with *AaArgM3* using the seed region complementarity and minimum

free energy (mfe) of -21 kcal/mol as the two main criteria.

To experimentally confirm the interaction between miRNAs and the target gene, *AaArgM3*, fragments of 200-500 bp long of *AaArgM3* 3'UTR containing the target sequences of aae-miR-2940, aae-miR-278, aae-miR-315, and aae-miR-1000 were amplified using primers with specific restriction sites XbaI and SacII. The fragments were then extracted from agarose gel, digested with XbaI and SacII, and ligated into pIZ/V5-His vector (Invitrogen) downstream of the *GFP* open reading frame. The right plasmids, confirmed by sequencing, were subsequently co-transfected into Sf9 cells (derived from *Spodoptera frugiperda*) together with control or miRNA mimics, respectively. All mimics were synthesized by Genepharma and used in transfection studies at a concentration of 100 μ M/ml. Cells were collected at 72 h after transfections, total RNA was extracted and RT-qPCR analyses were performed to determine the expression levels of the *GFP* gene. Three biological replicates with three technical replicates were analysed.

2.6. RNAi-mediated gene silencing

For RNAi-based experiments, dsRNAs were synthesized *in vitro* using the T7 Megascript transcription kit according to the manufacturer's instruction (Ambion Inc., USA). T7 promoter sequences (TAATACGACTCACTATAGGG) were incorporated in both forward and reverse primers designed to amplify a ~500 bp fragment of the *Ae. aegypti Dicer-1* (forward: 5'-CCCGGACCAAGTCCTAGTA-3'; reverse: 5'-CAACTCTTTCGGCACGTAA-3'), *AaArgM3* (forward: 5'-ATGCTATCCTCGATAACG-3'; reverse: 5'-TGCTATGATGTTAGCATTG-3') and the jellyfish *GFP* genes. For dsRNA synthesis, 200-500 ng of PCR product was used for each reaction. Reactions were incubated for 12 h at 37 °C, DNase-treated and precipitated by the lithium chloride method following the manufacturer's instructions. A total of 5 μ g of dsRNA was used to transfect Aag2 or aag2.wMelPop-CLA cells with 5 μ l of Cellfectin transfection reagent (Invitrogen). To reinforce silencing, cells were transfected again with the same reagent at 48 h after the first transfection. Cells were collected for RNA or DNA isolation as required for further analysis at 24 h after the second transfection. Gene silencing was confirmed by RT-qPCR using gene-

specific primers to *Dicer-1* and *AaArgM3* genes.

2.7. Western blotting

Cell samples were resuspended in PBS buffer to which 4×SDS-PAGE loading buffer was added. Proteins were separated on a 10% SDS-PAGE and transferred onto a nitrocellulose membrane. After blocking the membrane, it was probed with anti-GFP antibody (Abcam) and subsequently with alkaline phosphatase conjugated anti-rabbit antibody (Sigma). The same blot was subsequently probed with anti-histone H3 antibody (Invitrogen) to confirm equal loading of samples.

3. Results

3.1. Expression profile of *AaArgM3* in *Ae. aegypti* mosquito

By performing the NCBI BLAST, RNAHybrid and RNA22 software, a putative protein arginine methyltransferase 3 (*AaArgM3*, GeneBank ID: XM_001654962) from *Ae. aegypti* was identified as another target of a *Wolbachia* upregulated miRNA, aae-miR-2940-5p, which was previously confirmed to upregulate the transcript levels of the metalloprotease ftsh (*MetP*) gene (Hussain et al., 2011) and downregulate the transcript levels of *AaDnmt2* gene (Zhang et al., 2013). aae-miR-2940 is a mosquito-specific miRNA with its homolog absent in other insects (based on miRBase v.20). Sequence alignment showed that *AaArgM3* is a homologue of protein arginine methyltransferases, PRMT3 from human and DART3 from *Drosophila* (Bedford and Clarke, 2009; Boulanger et al., 2004). PRMT3 is a type I PRMT, and has been shown to be a cytosolic protein. Alignment results showed that there is 48% amino acid identity between *Drosophila* DART3 and *Ae. aegypti* *AaArgM3*.

In *Drosophila*, human and other animals, the expression of ArgM3 is developmentally and tissue-specifically regulated (Bedford and Clarke, 2009; Boulanger et al., 2004). By using *AaArgM3* gene-specific primers, we first investigated the expression pattern of *AaArgM3* in different developmental stages of *Ae. aegypti* by RT-PCR. Results showed that the transcripts of *AaArgM3* were detectable in the first and fourth instar larvae and adult female mosquitoes, but hardly

detectable in the second and third instar larvae (Fig. 1A). Further analysis showed that *AaArgM3* was mainly expressed in the abdomen of both male and female mosquitoes (Fig. 1B), which suggests that *AaArgM3* could be specifically expressed in some organs in the abdomen. Tissue-specific RT-qPCR analyses of five tissues (ovary, midgut, salivary, muscles and fatty body) from 4-day-old female *Ae. aegypti* confirmed that *AaArgM3* was mainly expressed in the ovary (Fig. 2A).

3.2. *Wolbachia* induces the expression of *AaArgM3* by using host miRNAs

It has been shown that *Wolbachia* manipulates host gene expression by regulating miRNA expression in *Ae. aegypti*, which improves colonization and blockage of DENV replication in the host (Bian et al., 2010; Hussain et al., 2011; Moreira et al., 2009; Osei-Amo et al., 2012; Zhang et al., 2013). Based on these, we investigated the expression of *AaArgM3* in female mosquitoes infected with *Wolbachia* and DENV using RT-qPCR. Results showed about two-fold higher transcript levels of *AaArgM3* in *Wolbachia*-infected mosquito tissues compared with those of the tet-cured mosquitoes (without *Wolbachia*; Wol⁻) (Fig. 2A-D). In *Ae. aegypti* mosquitoes infected with DENV, the transcript levels of *AaArgM3* did not significantly change compared with the mock-infected mosquitoes (Data not shown).

We also investigated the expression profiles of *AaArgM3* in *Ae. aegypti* cell lines infected with wMelPop-CLA (aag2.wMelPop-CLA) or without (Aag2) by RT-PCR. Results indicated that *AaArgM3* was expressed at much higher levels in aag2.wMelPop-CLA cells compared with Aag2 cells (Fig. 3A). To investigate whether miRNAs are involved in the regulation of *AaArgM3*, the Dicer-1 gene was silenced using RNAi in aag2.wMelPop-CLA cells. After confirming gene silencing, RT-PCR was carried out to explore the expression of *AaArgM3*. The expression levels of *AaArgM3* were considerably decreased compared with mock-transfected aag2.wMelPop-CLA cells (Fig. 3B), which suggested that the upregulation of *AaArgM3* expression in aag2.wMelPop-CLA cells could be mediated by miRNAs.

3.3. *AaArgM3* is targeted by aae-miR-2940

The target sequences of aae-miR-2940-5p were predicted in the 3' UTR of *AaArgM3* from

206 nucleotides 1991 to 2013 with significant complementarity to the miRNA's seed region (Fig. 4A).

207 To confirm the interaction of aae-miR-2940-5p with *AaArgM3*, we transfected aag2.wMelPop-CLA

208 cells with specific synthetic aae-miR-2940-5p and aae-miR-2940-3p inhibitors. RT-PCR results

209 showed much lower transcript levels of *AaArgM3* in the cells transfected with aae-miR-2940-5p

210 specific inhibitor compared with the cells transfected with the control aae-miR-2940-3p specific

211 inhibitor (Fig. 4B). To further validate the positive interaction of aae-miR-2940 with *AaArgM3*, the

212 target sequences were cloned downstream of the GFP gene in the pIZ/V5 vector (Fig. 5A). The

213 plasmid was subsequently co-transfected into Sf9 cells together with aae-miR-2940 mimic and a

214 control mimic (random sequences). The Sf9 cell line, which lacks the miRNA, provides an

215 independent system to test the miRNA-target interaction. RT-qPCR analyses were carried out to

216 assess the effect of miRNA-mRNA interaction on the transcript levels of the *GFP* gene. The results

217 showed that there were significantly higher levels of *GFP* transcripts in cells transfected with aae-

218 miR-2940 mimic compared to cells transfected with mock and control mimic (Fig. 5B). The

219 upregulation was also confirmed at the protein level using an anti-GFP antibody (Fig. 5C). These

220 results suggested that aae-miR-2940-5p upregulates the transcript levels of *AaArgM3*, which is

221 consistent with the expression pattern of *AaArgM3* gene in mosquitoes with or without *Wolbachia*

222 (Fig. 2).

223 Further bioinformatics analysis indicated that *AaArgM3* could also be a potential target of three

224 other miRNAs, aae-miR-278, -315, and -1000. We investigated the interaction of *AaArgM3* gene

225 with these predicted miRNAs by cloning their corresponding target sites in *AaArgM3* (Fig. 6A)

226 downstream of the *GFP* gene. The constructs were co-transfected into Sf9 cells with their

227 corresponding mimics. While aae-miR-278 and -1000 had no effect, aae-miR-315 mimic increased

228 *GFP* transcript levels compared with mock and the control mimic (Fig. 6B). However, when we

229 checked our previous microarray data (Hussain et al., 2011), we found that aae-miR-315 levels

230 slightly increased in *Wolbachia*-infected female mosquitoes but the difference was not significant.

231 Aae-miR-315 may regulate *AaArgM3* but perhaps not in the context of *Wolbachia*-mosquito

interaction.

3.4. *AaArgM3 facilitates Wolbachia replication*

In previous studies, we reported that *Wolbachia* infection leads to up- or downregulation of a number of host genes, which facilitate *Wolbachia* replication and maintenance in mosquito cells (Hussain et al., 2011; Osei-Amo et al., 2012; Zhang et al., 2013). Considering that aae-miR-2940-5p upregulates the transcript levels of *AaArgM3*, we first investigated whether *AaArgM3* has any effect on *Wolbachia* replication in aag2.wMelPop-CLA cells. For this, *AaArgM3* was silenced in the cells and the density of *Wolbachia* was analysed by qPCR. RT-qPCR confirmed that the silencing efficiency was over 90% (Fig. 7A). qPCR results with *wsp* gene-specific primers revealed that *Wolbachia* density was significantly lower in *AaArgM3* silenced cells, when compared with cells transfected with dsGFP or mock (Fig. 7B). This result suggests that *AaArgM3* enhances *Wolbachia* replication in the cell line, which is consistent with the expression profile that *AaArgM3* expression is considerably higher in the female mosquitoes with *Wolbachia*, compared with tet-cured counterpart mosquitoes (Fig. 2).

3.5. *AaArgM3 does not regulate DENV-2 replication*

In both *Ae. aegypti* mosquitoes and cell lines, *Wolbachia* was found to limit replication of DENV (Bian et al., 2010; Moreira et al., 2009), which could be due to manipulation of the host gene expression via miRNAs by *Wolbachia*. To explore the effect of *AaArgM3* on DENV replication, we silenced *AaArgM3* by transfecting Aag2 cells with *AaArgM3* dsRNA that were subsequently infected with DENV-2. Total RNA at 72 h after viral infection was isolated and analysed by RT-qPCR with DENV-specific primers. RT-qPCR confirmed that the silencing efficiency was about 85% (Fig. 8A). The results showed that the relative abundance of DENV was not significantly different in *AaArgM3* silenced cells compared with cells transfected with dsGFP or mock (Fig. 8B). Even silencing of the gene in aag2.wMelPop-CLA cells in which higher levels of *AaArgM3* are found, DENV replication was not different in mock, dsGFP or dsAaArgM3 cells (Fig. 8C). These results suggest that *AaArgM3* might not regulate replication of DENV in the mosquito cells, which

is consistent with the expression profile that the transcript levels of *AaArgM3* were not different in non-infected and DENV-infected mosquitoes.

4. Discussion

Utilization of *Wolbachia* has appeared as a viable non-chemical control strategy to limit transmission of vector-borne pathogens since they block replication of a variety of pathogens, including arboviruses. *Wolbachia* strains have been successfully introduced into *Ae. aegypti* and *An. gambiae*, the important vectors of dengue fever and malaria, and others in an effort to suppress transmission of DENV and *Plasmodium* (Bian et al., 2013; Bian et al., 2010; Blagrove et al., 2012; McMeniman et al., 2009; Xi et al., 2005). To survive and persist in the new hosts, the endosymbiotic bacteria have to evade or overcome host immune responses. Hussain et al. (2011) have previously reported that *Wolbachia* wMelPop-CLA strain induces differential expression of a number of host miRNAs, including the mosquito-specific aae-miR-2940, in *Ae. aegypti*. In *Ae. aegypti*, aae-miR-2940 upregulates the expression of one target gene, metalloprotease ftsh (*MetP*), which is crucial for efficient replication and maintenance of the endosymbiont (Hussain et al., 2011). Osei-Amo et al. (2012) found that differentially expressed aae-miR-12 downregulates the expression of two target genes, *MCT1* and *MCM6*, which also play a role in *Wolbachia*'s fitness in the mosquito cells. In addition, the methyltransferase *AaDnmt2* was identified to be another target of aae-miR-2940 and plays an important role in the replication of *Wolbachia* and contributes to the inhibition of DENV replication in *Ae. aegypti* (Zhang et al., 2013). These findings have shed light on molecular mechanisms by which *Wolbachia* manipulate the host's environment in *Ae. aegypti*. In the present study, *AaArgM3* was identified as another target gene of aae-miR-2940. The interaction of aae-miR-2940 with *AaArgM3* was confirmed and validated by using a synthetic inhibitor and mimic of aae-miR-2940 (Fig. 4B and 5). By examining the expression patterns, we found that the transcript levels of *AaArgM3* were significantly higher in *Wolbachia*-infected female mosquitoes (Fig. 2) and cells (Fig. 3A). Silencing of *AaArgM3* gene in aag2-wMelPop-CLA by RNAi showed a significant decline in *Wolbachia* density, but no effect on DENV (Fig. 7B, 8C).

284 Further, silencing of *AaArgM3* gene followed by DENV infection in Aag2 cells showed no
 285 significant effect on DENV replication. These results suggest that by inducing the expression of
 286 aae-miR-2940, *Wolbachia* upregulates the expression of *AaArgM3*, which in turn benefits
 287 *Wolbachia* in *Ae. aegypti*.

288 Methylation of arginine residues is a widespread posttranslational modification of proteins
 289 catalyzed by a conserved family of protein arginine methyltransferases. Protein arginine
 290 methyltransferases are classified into three types by methylated arginine residues including
 291 asymmetric ω - N^G , N^G -dimethylarginine (ADMA), symmetric ω - N^G , N^G -dimethylarginine (SDMA)
 292 and ω - N^G -dimethylarginine (MMA). Type I includes PRMT1, 2, 3, 4 and 8; type II includes
 293 PRMT5 and 7 and type III includes PRMT7 (Bedford and Clarke, 2009; McBride and Silver, 2001).
 294 They have diverse biological roles in the regulation of a large array of cell processes including
 295 signal transduction, subcellular localization, RNA processing and transcription (Bedford and
 296 Clarke, 2009; Krause et al., 2007; McBride and Silver, 2001). In recent years, PRMTs from
 297 mammals have been found to play essential roles in regulating the replication, production and
 298 infectivity of a variety of viruses. For example, PRMT1 negatively regulated Hepatitis Delta virus
 299 (Li et al., 2004), hepatitis B virus (Benhenda et al., 2013), hepatitis C virus (Duong et al., 2005) and
 300 Herpes Simplex virus (Souki et al., 2009; Yu et al., 2010). PRMT1 and PRMT5 together repressed
 301 HIV long terminal repeat transcription and consequently suppressed replication of the virus (Kwak
 302 et al., 2003). PRMT6 inhibited HIV-1 transcription through the methylation of Tat, Rev and
 303 nucleocapsid proteins (Boulanger et al., 2005; Invernizzi et al., 2007; Invernizzi et al., 2006;
 304 Singhroy et al., 2013; Xie et al., 2007). In our preliminary experiment, exposure of Aag2 cells to a
 305 protein arginine methyltransferase inhibitor (adenosine-2,3-dialdehyde) led to increased DENV
 306 replication (Zhang et al., unpublished data). In this study, we did not find that silencing of
 307 *AaArgM3* had any effect on DENV replication, but we cannot exclude the possible role of
 308 *AaArgM3* in regulating DENV replication. This is because in *Ae. aegypti* there are eight family
 309 members of protein arginine methyltransferases, which could have overlapping function probably

compensating the function of AaArgM3 when it was silenced. Further study is required to investigate which family member(s) play a role in regulating DENV replication.

miRNAs have been implicated as gene regulators controlling diverse biological processes including development, cancer, immunity and host–microorganism interactions. They usually downregulate their target genes by either degradation of the target mRNA or repression of translation (reviewed in Asgari, 2013; Bartel, 2009). A large number of miRNAs have been identified to control the DNA and RNA methylation machineries (Denis et al., 2011). However, very few miRNAs have been identified to regulate protein arginine methylation. Recently miR-181a, b, c, and d family members were found to directly regulate *CARM1* (*PRMT4*) expression in human embryonic stem cells (Xu et al., 2013). All the miR-181 family members target the 3' UTR of *CARM1*.

In our study, we identified and confirmed that aae-miR-2940, which is induced in the presence of *Wolbachia*, enhances the expression of a protein arginine methyltransferase, *AaArgM3*, in *Ae. aegypti*, which appears to be important for *Wolbachia* fitness. This suggests a positive feedback loop in which *Wolbachia* infection induces aae-miR-2940 that in turn positively regulates *AaArgM3* leading to more *Wolbachia*. However, the mechanism by which the protein facilitates *Wolbachia* maintenance remains to be investigated. Our results suggest that *Wolbachia* manipulates host physiology and gene expression for colonization in mosquitoes using multiple targets of differentially regulated miRNAs.

Acknowledgement

The authors would like to acknowledge the financial support from the Australian Research Council Discovery grant to SA (DP110102112) and an ARC DECRA fellowship to MH (DE120101512).

References

- Asgari, S., 2013. MicroRNA functions in insects. *Insect Biochem Mol Biol* 43, 388-397.
- Baldini, F., Segata, N., Pompon, J., Marcenac, P., Robert Shaw, W., Dabiré, R., Diabaté, A., Levashina, E., Catteruccia, F., 2014. Evidence of natural *Wolbachia* infections in field populations of *Anopheles gambiae*. *Nat Commun* 5, 3985.

- 336 Bartel, D.P., 2009. MicroRNAs: target recognition and regulatory functions. *Cell* 23, 215-233.
- 337 Bedford, M.T., Clarke, S.G., 2009. Protein arginine methylation in mammals: Who, what, and why.
338 *Mol. Cell* 33, 1-13.
- 339 Benhenda, S., Ducroux, A., Riviere, L., Sobhian, B., Ward, M.D., Dion, S., Hantz, O., Protzer, U.,
340 Michel, M.L., Benkirane, M., Semmes, O.J., Buendia, M.A., Neuveut, C., 2013.
341 Methyltransferase PRMT1 Is a binding partner of HBx and a negative regulator of Hepatitis B
342 virus transcription. *J Virol* 87, 4360-4371.
- 343 Bian, G., Joshi, D., Dong, Y., Lu, P., Zhou, G., Pan, X., Xu, Y., Dimopoulos, G., Xi, Z., 2013.
344 *Wolbachia* invades *Anopheles stephensi* populations and induces refractoriness to *Plasmodium*
345 infection. *Science* 340, 748-751.
- 346 Bian, G., Xu, Y., Lu, P., Xie, Y., Xi, Z., 2010. The endosymbiotic bacterium *Wolbachia* induces
347 resistance to Dengue virus in *Aedes aegypti*. *PLoS Pathog* 6, e1000833.
- 348 Blagrove, M.S.C., Arias-Goeta, C., Failloux, A.-B., Sinkins, S.P., 2012. *Wolbachia* strain wMel
349 induces cytoplasmic incompatibility and blocks dengue transmission in *Aedes albopictus*. *Proc*
350 *Natl Acad Sci USA* 102, 255-260.
- 351 Boulanger, M.C., Liang, C., Russell, R.S., Lin, R.T., Bedford, M.T., Wainberg, M.A., Richard, S.,
352 2005. Methylation of tat by PRMT6 regulates human immunodeficiency virus type 1 gene
353 expression. *J Virol* 79, 124-131.
- 354 Boulanger, M.C., Miranda, T.B., Clarke, S., di Fruscio, M., Suter, B., Lasko, P., Richard, S., 2004.
355 Characterization of the *Drosophila* protein arginine methyltransferases DART1 and DART4.
356 *Biochem J* 379, 283-289.
- 357 Bourtzis, K., Pettigrew, M., O'Neill, S., 2000. *Wolbachia* neither induces nor suppresses transcripts
358 encoding antimicrobial peptides. *Insect Mol Biol* 9, 635-639.
- 359 Denis, H., Ndlovu, M.N., Fuks, F., 2011. Regulation of mammalian DNA methyltransferases: a
360 route to new mechanisms. *EMBO Rep* 12, 647-656.

- 361 Duong, F.H.T., Christen, V., Berke, J.M., Penna, S.H., Moradpour, D., Heim, M.H., 2005.
362 Upregulation of protein phosphatase 2Ac by hepatitis C virus modulates NS3 helicase activity
363 through inhibition of protein arginine methyltransferase 1. J Virol 79, 15342-15350.
- 364 Fehri, L.F., Koch, M., Belogolova, E., Khalil, H., Bolz, C., Kalali, B., Mollenkopf, H.J., Beigier-
365 Bompadre, M., Karlas, A., Schneider, T., Churin, Y., Gerhard, M., Meyer, T.F., 2010.
366 *Helicobacter pylori* induces miR-155 in T cells in a cAMP-Foxp3-dependent manner. PLoS
367 ONE 5, e9500.
- 368 Frentiu, F.D., Robinson, J., Young, P.R., McGraw, E.A., O'Neill, S.A., 2010. *Wolbachia*-mediated
369 resistance to Dengue virus infection and death at the cellular level. PLoS ONE 5, e13398.
- 370 Frentiu, F.D., Zakir, T., Walker, T., Popovici, J., Pyke, A.T., van den Hurk, A., McGraw, E.A.,
371 O'Neill, S.L., 2014. Limited dengue virus replication in field-collected *Aedes aegypti*
372 mosquitoes infected with *Wolbachia*. PLoS Negl Trop Dis 8, e2688.
- 373 Hilgenboecker, K., Hammerstein, P., Schlattmann, P., Telschow, A., Werren, J.H., 2008. How
374 many species are infected with *Wolbachia*?—A statistical analysis of current data. FEMS
375 Microbiol Lett 218, 215–220.
- 376 Huang, J., Wang, F., Argyris, E., Chen, K., Liang, Z., Tian, H., Huang, W., Squires, K.,
377 Verlinghieri, G., Zhang, H., 2007. Cellular microRNAs contribute to HIV-1 latency in resting
378 primary CD4+ T lymphocytes. Nat Med 13, 1241-1247.
- 379 Hussain, M., Frentiu, F.D., Moreira, L.A., O'Neill, S.L., Asgari, S., 2011. *Wolbachia* utilizes host
380 microRNAs to manipulate host gene expression and facilitate colonization of the dengue vector
381 *Aedes aegypti*. Proc Natl Acad Sci USA 108, 9250-9255.
- 382 Invernizzi, C.F., Xie, B.D., Frankel, F.A., Feldhammer, M., Roy, B.R., Richard, S., Wainberg,
383 M.A., 2007. Arginine methylation of the HIV-1 nucleocapsid protein results in its diminished
384 function. AIDS 21, 795-805.
- 385 Invernizzi, C.F., Xie, B.D., Richard, S., Wainberg, M.A., 2006. PRMT6 diminishes HIV-1 Rev
386 binding to and export of viral RNA. Retrovirology 3, 93.

- 387 Jeyaparakash, A., Hoy, M.A., 2000. Long PCR improves *Wolbachia* DNA amplification: wsp
388 sequences found in 76% of sixty-three arthropod species. *Insect Mol Biol* 9, 393-405.
- 389 Kambris, Z., Blagborough, A.M., Pinto, S.B., Blagrove, M.S.C., Godfray, H.C.J., Sinden, R.E.,
390 Sinkins, S.P., 2010. *Wolbachia* stimulates immune gene expression and inhibits *Plasmodium*
391 development in *Anopheles gambiae*. *PLoS Pathog* 6, e1001143.
- 392 Kambris, Z., Cook, P.E., Phuc, H.K., Sinkins, S.P., 2009. Immune activation by life-shortening
393 *Wolbachia* and reduced filarial competence in mosquitoes. *Science* 326, 134-136.
- 394 Krause, C., Yang, Z., Kim, Y., Lee, J., Cook, J., Pestka, S., 2007. Protein arginine
395 methyltransferases: evolution and assessment of their pharmacological and therapeutic
396 potential. *Pharmacol Ther* 113, 50-87.
- 397 Kwak, Y.T., Guo, J., Prajapati, S., Park, K.J., Surabhi, R.M., Miller, B., Gehrig, P., Gaynor, R.B.,
398 2003. Methylation of SPT5 regulates its interaction with RNA polymerase II and transcriptional
399 elongation properties. *Mol Cell* 11, 1055-1066.
- 400 Li, Y.J., Stallcup, M.R., Lai, M.M.C., 2004. Hepatitis delta virus antigen is methylated at arginine
401 residues, and methylation regulates subcellular localization and RNA replication. *J Virol* 78,
402 13325-13334.
- 403 Lu, F., Weidmer, A., Liu, C.G., Volinia, S., Croce, C.M., Lieberman, P.M., 2008. Epstein-Barr
404 virus-induced miR-155 attenuates NF-kappa B signaling and stabilizes latent virus persistence.
405 *J Virol* 82, 10436-10443.
- 406 McBride, A.E., Silver, P.A., 2001. State of the Arg: Protein methylation at arginine comes of age.
407 *Cell* 106, 5-8.
- 408 McMeniman, C.J., Lane, R.V., Cass, B.N., Fong, A.W.C., Sidhu, M., Wang, Y.-F., O'Neill, S.L.,
409 2009. Stable introduction of a life-shortening *Wolbachia* infection into the mosquito *Aedes*
410 *aegypti*. *Science* 323, 141-144.
- 411 Moreira, L.A., Iturbe-Ormaetxe, I., Jeffery, J.A., Lu, G.J., Pyke, A.T., Hedges, L.M., Rocha, B.C.,
412 Hall-Mendelin, S., Day, A., Riegler, M., Hugo, L.E., Johnson, K.N., Kay, B.H., McGraw, E.A.,

- 413 van den Hurk, A.F., Ryan, P.A., O'Neill, S.L., 2009. A *Wolbachia* symbiont in *Aedes aegypti*
414 limits infection with Dengue, Chikungunya, and *Plasmodium*. *Cell* 139, 1268-1278.
- 415 Osei-Amo, S., Hussain, M., O'Neill, S.L., Asgari, S., 2012. *Wolbachia*-induced aae-miR-12
416 miRNA negatively regulates the expression of MCT1 and MCM6 genes in *Wolbachia*-infected
417 mosquito cell line. *PLoS ONE* 7, e50049.
- 418 Pan, X., Zhou, G., Wu, J., Bian, G., Lu, P., Raikhel, A.S., Xi, Z., 2012. *Wolbachia* induces reactive
419 oxygen species (ROS)-dependent activation of the Toll pathway to control dengue virus in the
420 mosquito *Aedes aegypti*. *Proc Natl Acad Sci USA* 109, E23-31.
- 421 Rainey, S., Shah, P., Kohl, A., Dietrich, I., 2014. Understanding the *Wolbachia*-mediated inhibition
422 of arboviruses in mosquitoes: progress and challenges. *J Gen Virol* 95, 517-530.
- 423 Rances, E., Johnson, T.K., Popovici, J., Iturbe-Ormaetxe, I., Zakir, T., Warr, C.G., O'Neill, S.L.,
424 2013. The Toll and Imd pathways are not required for *Wolbachia*-mediated dengue interference.
425 *J Virol* 8, e1002548.
- 426 Rancès, E., Ye, Y.H., Woolfit, M., McGraw, E.A., O'Neill, S.L., 2012. The relative importance of
427 innate immune priming in *Wolbachia*-mediated dengue interference. *PLoS Pathog* 8, e1002548.
- 428 Rehmsmeier, M., Steffen, P., Höchsmann, M., Giegerich, R., 2004. Fast and effective prediction of
429 microRNA/target duplexes. *RNA* 10, 1507-1517.
- 430 Singhroy, D.N., Mesplede, T., Sabbah, A., Quashie, P.K., Falgout, J.P., Wainberg, M.A., 2013.
431 Automethylation of protein arginine methyltransferase 6 (PRMT6) regulates its stability and its
432 anti-HIV-1 activity. *Retrovirology* 10, 73.
- 433 Souki, S.K., Gershon, P.D., Sandri-Goldin, R.M., 2009. Arginine methylation of the ICP27 RGG
434 box regulates ICP27 export and is required for efficient Herpes Simplex virus 1 replication. *J*
435 *Virol* 83, 5309-5320.
- 436 Tili, E., Michaille, J.-J., Cimino, A., Costinean, S., Dumitru, C.D., Adair, B., Fabbri, M., Alder, H.,
437 Liu, C.G., Calin, G.A., Croce, C.M., 2007. Modulation of miR-155 and miR-125b levels

following lipopolysaccharide/TNF- α stimulation and their possible roles in regulating the response to endotoxin shock. J Immunol 179, 5082-5089.

Walker, T., Johnson, P.H., Moreira, L.A., Iturbe-Ormaetxe, I., Frentiu, F.D., McMeniman, C.J., Leong, Y.S., Dong, Y., Axford, J., Kriesner, P., Lloyd, A.L., Ritchie, S.A., O'Neill, S.L., Hoffman, A.A., 2011. The wMel *Wolbachia* strain blocks dengue and invades caged *Aedes aegypti* populations. Nature 476, 450-453.

Xi, Z., Gavotte, L., Xie, Y., Dobson, S.L., 2008. Genome-wide analysis of the interaction between the endosymbiotic bacterium *Wolbachia* and its *Drosophila* host. BMC Genomics 9, 1.

Xi, Z., Khoo, C., Dobson, S., 2005. *Wolbachia* establishment and invasion in an *Aedes aegypti* laboratory population. Science 310, 326-328.

Xie, B., Invernizzi, C.F., Richard, S., Wainberg, M.A., 2007. Arginine methylation of the human immunodeficiency virus type 1 Tat protein by PRMT6 negatively affects Tat interactions with both cyclin T1 and the Tat transactivation region. J Virol 81, 4226-4234.

Xu, Z.Y., Jiang, J.F., Xu, C., Wang, Y., Sun, L., Guo, X.C., Liu, H.Q., 2013. MicroRNA-181 regulates CARM1 and histone arginine methylation to promote differentiation of human embryonic stem cells. PLoS ONE 8, e53146.

Yu, J., Shin, B., Park, E.S., Yang, S., Choi, S., Kang, M., Rho, J., 2010. Protein arginine methyltransferase 1 regulates herpes simplex virus replication through ICP27 RGG-box methylation. Biochem Biophys Res Commun 391, 322-328.

Zhang, G., Hussain, M., O'Neill, S.L., Asgari, S., 2013. *Wolbachia* uses a host microRNA to regulate transcripts of a methyltransferase contributing to dengue virus inhibition in *Aedes aegypti*. Proc Natl Acad Sci USA 110, 10276-10281.

Zug, R., Hammerstein, P., 2012. Still a host of hosts for *Wolbachia*: analysis of recent data suggests that 40% of terrestrial arthropod species are infected. PLoS ONE 7, e38544.

Figure Legends

Fig. 1. AaArgM3 expression in *Ae. aegypti*. (A) RT-PCR analysis was performed using the total RNA samples from *Ae. aegypti* mosquito larvae and female adults. (B) RT-PCR analysis was performed with head+thorax (HT) and abdomen (Ab) of mosquito females and males. *Rps17* gene was used as control to show the integrity of RNA.

Fig. 2. Tissue-specific expression of AaArgM3 in female *Ae. aegypti* mosquitoes. RT-qPCR analysis of transcript levels of *AaArgM3* in ovary, midgut, salivary glands (Salivary), thoracic muscle tissues (Muscle) and fat body from 4-day-old (A) tetracycline-treated non-infected (Wol⁻) and (B) *Wolbachia*-infected (Wol⁺) female mosquitoes. The transcript levels of *AaArgM3* were also compared in (C) the ovaries and (D) fat body in the samples. Asterisks indicate a significant difference between treatments (** $p < 0.001$).

Fig. 3. AaArgM3 expression in aag2-wMelPop-CLA and Aag2 cells. (A) RT-PCR analysis of RNA extracted from aag2-wMelPop-CLA (Pop) and Aag2 cells. (B) RT-PCR analysis of RNA extracted from mock and dsDicer-1 transfected aag2-wMelPop-CLA cells. *Rps17* gene was used as control to show the integrity of RNA.

Fig. 4. AaArgM3 transcript levels are upregulated by aae-miR-2940-5p. (A) Schematic diagram showing the *AaArgM3* mRNA and its target sequences with complete complementarity of aae-miR-2940-5p seed region (bold and underlined) with the sequences. (B) RT-PCR analysis of *AaArgM3* relative transcript levels using RNA extracted from aag2.wMelPop-CLA cells transfected with mock, synthetic aae-miR-2940-5p or aae-miR-2940-3p (control) inhibitors. *Rps17* gene was used as control to show the integrity of RNA.

Fig. 5. Target validation of aae-miR-2940. (A) Schematic diagram showing the cloning strategy of *AaArgM3* target sequence complementary to the miRNA seed region from the *AaArgM3* 3'UTR under the *GFP* reporter gene in the pIZ vector. (B) RT-qPCR analysis of *GFP* transcript levels using the RNA extracted from Sf9 cells co-transfected with pIZ-GFP-target and mock, synthetic

control mimic or aae-miR-2940 mimic. *Actin* gene was used as the normalizing control. Asterisks indicate a significant difference between mock or control mimic and aae-miR-2940 mimic transfections ($p < 0.0001$). (C) Western blot analysis of Sf9 cells transfected with pIZ/GFP-target together with aae-2940-5p mimic (2940), control mimic (Cmimic), no mimic (Nmimic) or mock transfected without plasmid (Mock). The blot was probed with anti-GFP antibody and subsequently with anti-histone H3 to show equal loading of samples.

Fig. 6. Interactions of *AaArgM3* with predicted miRNAs. (A) *Ae. aegypti* *AaArgM3* was predicted to be the target of aae-miR-278, -315, and -1000 with complete complementarity of their seed regions (bold and underlined) with the sequences. (B) RT-qPCR analysis of GFP expression using RNA extracted from Sf9 cells transfected with pIZ-GFP-target and mock, synthetic control mimic, aae-miR-278, -315 or -1000 mimics. *Actin* gene was used as the normalizing gene. There are statistically significant differences between treatments with different letters at $p < 0.05$.

Fig. 7. *AaArgM3* facilitates *Wolbachia* replication. RNAi-mediated silencing of *AaArgM3* gene was carried out in aag2.wMelPop-CLA cells for 72 h. (A) RT-qPCR analysis of *AaArgM3* gene relative to *RPS17* in aag2.wMelPop-CLA cells transfected with mock, GFP and *AaArgM3* dsRNAs. (B) qPCR analysis of *Wolbachia* density in aag2.wMelPop-CLA cells 72 h after transfection with mock, GFP and *AaArgM3* dsRNAs using primers specific to the *Wolbachia* *wsp* gene. Asterisks indicate a significant difference between transfection with *AaArgM3* dsRNA and other treatments (*** $p < 0.0001$; ** $p < 0.001$).

Fig. 8. *AaArgM3* has no effect on DENV replication in Aag2 or *Wolbachia*-infected Aag2 cells. RNAi-mediated silencing of *AaArgM3* gene was carried out in Aag2 cells. 72 h after transfection with dsRNA, cells were infected with DENV-2. At 72 h after infection, total RNA was extracted from cells. (A) RT-qPCR analysis of *AaArgM3* gene relative to *RPS17* in Aag2 cells transfected with mock, GFP and *AaArgM3* dsRNAs and infected with DENV-2. (B) RT-qPCR analysis of RNA using DENV-specific primers in Aag2 cells transfected with mock, GFP and *AaArgM3*

513 dsRNAs and then infected with DENV-2. (C) RT-qPCR analysis of RNA using DENV-specific
514 primers from aag2.wMelPop-CLA cells transfected with mock, GFP and *AaArgM3* dsRNAs and
515 then infected with DENV-2 for 72 h. Silencing of *AaArgM3* in aag2.wMelPop-CLA cells was
516 confirmed as shown in Fig. 7A. Asterisks indicate a significant difference between transfection with
517 *AaArgM3* dsRNA and other treatments (***) $p < 0.0001$).

ACCEPTED MANUSCRIPT


Figure 1

Figure 2





Figure 3

Figure 4




Figure 5




Figure 6**A****AaArgM3/miR278 predicted interaction at 588-609**

```

target 5' C          UUACU          U 3'
          GAAUGGAUGGG      UCCUGCUG
          UUUGCCUGCUU      AGGGUGGC
miRNA  3'          UC          U 5'

```

AaArgM3/miR315 predicted interaction at 601-622

```

target 5' U      CC          G 3'
          ACUU      UGCUGUUCGAGGG
          UGAA      AUGACGAGCUUUC
miRNA  3' C      ACUA          5'

```

AaArgM3/miR1000 predicted interaction at 606-627

```

target 5' C          UC  G          G 3'
          CUGCUGU  GA  GGAUGAUG
          GACGACA  CU  UCCUGUUUAU
miRNA  3' U          G          A 5'

```


B

Figure 7





Figure 8



Highlights

- Arginine methyltransferase was found as another target of aae-miR-2940-5p, a mosquito-specific miRNA
- Arginine methyltransferase is induced in *Wolbachia*-infected *Aedes aegypti* mosquitoes and cells
- Arginine methyltransferase is positively regulated by aae-miR-2940-5p.
- Arginine methyltransferase contributes to replication/maintenance of *Wolbachia* but has no effect on dengue virus replication.